

Quotients of Tannakian Categories

J.S. Milne*

August 24, 2005; July 9, 2007

Abstract

We classify the “quotients” of a tannakian category in which the objects of a tannakian subcategory become trivial, and we examine the properties of such quotient categories.

Introduction

Given a tannakian category T and a tannakian subcategory S , we ask whether there exists a quotient of T by S , by which we mean an exact tensor functor $q: T \rightarrow Q$ from T to a tannakian category Q such that

- (a) the objects of T that become trivial in Q (i.e., isomorphic to a direct sum of copies of $\mathbb{1}$ in Q) are precisely those in S , and
- (b) every object of Q is a subquotient of an object in the image of q .

When T is the category $\text{Rep}(G)$ of finite-dimensional representations of an affine group scheme G the answer is obvious: there exists a unique normal subgroup H of G such that the objects of S are the representations on which H acts trivially, and there exists a canonical functor q satisfying (a) and (b), namely, the restriction functor $\text{Rep}(G) \rightarrow \text{Rep}(H)$ corresponding to the inclusion $H \hookrightarrow G$. By contrast, in the general case, there need not exist a quotient, and when there does there will usually not be a canonical one. In fact, we prove that there exists a q satisfying (a) and (b) if and only if S is neutral, in which case the q are classified by the k -valued fibre functors on S . Here $k \stackrel{\text{def}}{=} \text{End}(\mathbb{1})$ is assumed to be a field.

From a slightly different perspective, one can ask the following question: given a subgroup H of the fundamental group $\pi(T)$ of T , does there exist an exact tensor functor $q: T \rightarrow Q$ such that the resulting homomorphism $\pi(Q) \rightarrow q(\pi(T))$ maps $\pi(Q)$ isomorphically onto $q(H)$? Again, there exists such a q if and only if the subcategory T^H of T , whose objects are those on which H acts trivially, is neutral, in which case the functors q correspond to the k -valued fibre functors on T^H .

The two questions are related by the “tannakian correspondence” between tannakian subcategories of T and subgroups of $\pi(T)$ (see 1.7).

* Available at www.jmilne.org/math/.

In addition to proving the above results, we determine the fibre functors, polarizations, and fundamental groups of the quotient categories $\mathcal{Q}$.

The original motivation for these investigations came from the theory of motives (see Milne 2002, 2007).

Notation: The notation $X \approx Y$ means that X and Y are isomorphic, and $X \simeq Y$ means that X and Y are canonically isomorphic (or that there is a given or unique isomorphism).

1 Preliminaries

For tannakian categories, we use the terminology of Deligne and Milne 1982. In particular, we write $\mathbb{1}$ for any identity object of a tannakian category — recall that it is uniquely determined up to a unique isomorphism. We fix a field k and consider only tannakian categories with $k = \text{End}(\mathbb{1})$ and only functors of tannakian categories that are k -linear.

Gerbes

1.1 We refer to Giraud 1971, Chapitre IV, for the theory of gerbes. All gerbes will be for the flat (i.e., fpqc) topology on the category Aff_k of affine schemes over k . The band (= lien) of a gerbe $\mathcal{G}$ is denoted $\text{Bd}(\mathcal{G})$. A commutative band can be identified with a sheaf of groups.

1.2 Let $\alpha: \mathcal{G}_1 \rightarrow \mathcal{G}_2$ be a morphism of gerbes over Aff_k , and let ω_0 be an object of $\mathcal{G}_{2,k}$. Define $(\omega_0 \downarrow \mathcal{G}_1)$ to be the fibred category over Aff_k whose fibre over $S \xrightarrow{s} \text{Spec } k$ has as objects the pairs (ω, a) consisting of an object ω of $\text{ob}(\mathcal{G}_{1,S})$ and an isomorphism $a: s^*\omega_0 \rightarrow \alpha(\omega)$ in $\mathcal{G}_{2,S}$; the morphisms $(\omega, a) \rightarrow (\nu, b)$ are the isomorphisms $\varphi: \omega \rightarrow \nu$ in $\mathcal{G}_{1,S}$ giving rise to a commutative triangle. Thus,

$$\begin{array}{ccc}
 \omega & & \alpha(\omega) \\
 \downarrow \varphi & \nearrow a & \downarrow \alpha(\varphi) \\
 \nu & s^*(\omega_0) & \alpha(\nu) \\
 & \searrow b & \\
 \mathcal{G}_{1,S} & & \mathcal{G}_{2,S}
 \end{array}$$

If the map of bands defined by α is an epimorphism, then $(\omega_0 \downarrow \mathcal{G}_1)$ is a gerbe, and the sequence of bands

$$1 \rightarrow \text{Bd}(\omega_0 \downarrow \mathcal{G}_1) \rightarrow \text{Bd}(\mathcal{G}_1) \rightarrow \text{Bd}(\mathcal{G}_2) \rightarrow 1 \quad (1)$$

is exact (Giraud 1971, IV 2.5.5(i)).

1.3 Recall (Saavedra Rivano 1972, III 2.2.2) that a gerbe is said to be tannakian if its band is locally defined by an affine group scheme. It is clear from the exact sequence (1) that if $\mathcal{G}_1$ and $\mathcal{G}_2$ are tannakian, then so also is $(\omega_0 \downarrow \mathcal{G}_1)$.

1.4 The fibre functors on a tannakian category $\mathcal{T}$ form a gerbe $\text{FIB}(\mathcal{T})$ over Aff_k (Deligne 1990, 1.13). Each object X of $\mathcal{T}$ defines a representation $\omega \mapsto \omega(X)$ of $\text{FIB}(\mathcal{T})$, and in this way we get an equivalence $\mathcal{T} \rightarrow \text{Rep}(\text{FIB}(\mathcal{T}))$ of tannakian categories (Deligne 1989, 5.11; Saavedra Rivano 1972, III 3.2.3, p200). Every gerbe whose band is tannakian arises in this way from a tannakian category (Saavedra Rivano 1972, III 2.2.3).

Fundamental groups

1.5 We refer to Deligne 1989, §§5,6, for the theory of algebraic geometry in a tannakian category $\mathcal{T}$ and, in particular, for the fundamental group $\pi(\mathcal{T})$ of $\mathcal{T}$. It is the affine group scheme¹ in $\mathcal{T}$ such that $\omega(\pi(\mathcal{T})) \simeq \underline{\text{Aut}}^{\otimes}(\omega)$ functorially in the fibre functor ω on $\mathcal{T}$. The group $\pi(\mathcal{T})$ acts on each object X of $\mathcal{T}$, and ω transforms this action into the natural action of $\underline{\text{Aut}}^{\otimes}(\omega)$ on $\omega(X)$. The various realizations $\omega(\pi(\mathcal{T}))$ of $\pi(\mathcal{T})$ determine the band of $\mathcal{T}$ (i.e., the band of $\text{FIB}(\mathcal{T})$).

1.6 An exact tensor functor $F: \mathcal{T}_1 \rightarrow \mathcal{T}_2$ of tannakian categories defines a homomorphism $\pi(F): \pi(\mathcal{T}_2) \rightarrow F(\pi(\mathcal{T}_1))$ (Deligne 1989, 6.4). Moreover:

- (a) F induces an equivalence of $\mathcal{T}_1$ with a category whose objects are the objects of $\mathcal{T}_2$ endowed with an action of $F(\pi(\mathcal{T}_1))$ compatible with that of $\pi(\mathcal{T}_2)$ (Deligne 1989, 6.5);
- (b) $\pi(F)$ is flat and surjective if and only if F is fully faithful and every subobject of $F(X)$, for X in $\mathcal{T}_1$, is isomorphic to the image of a subobject of X (cf. Deligne and Milne 1982, 2.21);
- (c) $\pi(F)$ is a closed immersion if and only if every object of $\mathcal{T}_2$ is a subquotient of an object in the image of q (ibid.).

1.7 For a subgroup² $H \subset \pi(\mathcal{T})$, we let $\mathcal{T}^H$ denote the full subcategory of $\mathcal{T}$ whose objects are those on which H acts trivially. It is a tannakian subcategory of $\mathcal{T}$ (i.e., it is a strictly full subcategory closed under the formation of subquotients, direct sums, tensor products, and duals) and every tannakian subcategory arises in this way from a unique subgroup of $\pi(\mathcal{T})$ (cf. Bertolin 2003, 1.6). The objects of $\mathcal{T}^{\pi(\mathcal{T})}$ are exactly the trivial objects of $\mathcal{T}$, and there exists a unique (up to a unique isomorphism) fibre functor

$$\gamma^{\mathcal{T}}: \mathcal{T}^{\pi(\mathcal{T})} \rightarrow \text{Vec}_k,$$

namely, $\gamma^{\mathcal{T}}(X) = \text{Hom}(\mathbb{1}, X)$.

1.8 For a subgroup H of $\pi(\mathcal{T})$ and an object X of $\mathcal{T}$, we let X^H denote the largest subobject of X on which the action of H is trivial. Thus $X = X^H$ if and only if X is in $\mathcal{T}^H$.

¹“ $\mathcal{T}$ -schéma en groupes affines” in Deligne’s terminology.

²Note that every subgroup H of $\pi(\mathcal{T})$ is normal. For example, the fundamental group π of the category $\text{Rep}(G)$ of representations of the affine group scheme $G = \text{Spec}(A)$ is A regarded as an object of $\text{Ind}(\text{Rep}(G))$. The action of G on A is that defined by inner automorphisms. A subgroup of π is a quotient $A \rightarrow B$ of A (as a bi-algebra) such that the action of G on A defines an action of G on B . Such quotients correspond to normal subgroups of G .

1.9 When H is contained in the centre of $\pi(\mathcal{T})$, then it is an affine group scheme in $\mathcal{T}^{\pi(\mathcal{T})}$, and so $\gamma^{\mathcal{T}}$ identifies it with an affine group scheme over k in the usual sense. For example, $\gamma^{\mathcal{T}}$ identifies the centre of $\pi(\mathcal{T})$ with $\underline{\text{Aut}}^{\otimes}(\text{id}_{\mathcal{T}})$ (cf. Saavedra Rivano 1972, II 3.3.3.2, p150).

Morphisms of tannakian categories

1.10 For a group G , a right G -object X , and a left G -object Y , $X \wedge^G Y$ denotes the contracted product of X and Y , i.e., the quotient of $X \times Y$ by the diagonal action of G , $(x, y)g = (xg, g^{-1}y)$. When $G \rightarrow H$ is a homomorphism of groups, $X \wedge^G H$ is the H -object obtained from X by extension of the structure group. In this last case, if X is a G -torsor, then $X \wedge^G H$ is also an H -torsor. See Giraud 1971, III 1.3, 1.4.

1.11 Let $\mathcal{T}$ be a tannakian category over k , and assume that the fundamental group π of $\mathcal{T}$ is commutative. A torsor P under π in $\mathcal{T}$ defines a tensor equivalence $\mathcal{T} \rightarrow \mathcal{T}$, $X \mapsto P \wedge^{\pi} X$, bound by the identity map on $\text{Bd}(\mathcal{T})$, and every such equivalence arises in this way from a torsor under π (cf. Saavedra Rivano 1972, III 2.3). For any k -algebra R and R -valued fibre functor ω on $\mathcal{T}$, $\omega(P)$ is an R -torsor under $\omega(\pi)$ and $\omega(P \wedge^{\pi} X) \simeq \omega(P) \wedge^{\omega(\pi)} \omega(X)$.

2 Quotients

For any exact tensor functor $q: \mathcal{T} \rightarrow \mathcal{T}'$, the full subcategory $\mathcal{T}^q$ of $\mathcal{T}$ whose objects become trivial in $\mathcal{T}'$ is a tannakian subcategory of $\mathcal{T}$ (obviously).

We say that an exact tensor functor $q: \mathcal{T} \rightarrow \mathcal{Q}$ of tannakian categories is a **quotient functor** if every object of $\mathcal{Q}$ is a subquotient of an object in the image of q ; equivalently, if the homomorphism $\pi(q): \pi(\mathcal{Q}) \rightarrow q(\pi\mathcal{T})$ is a closed immersion (see 1.6(c)). If, in addition, the homomorphism $\pi(q)$ is normal (i.e., its image is a normal subgroup of $q(\mathcal{T})$), then we say that q is **normal**.

EXAMPLE 2.1 Consider the exact tensor functor $\omega^f: \text{Rep}(G) \rightarrow \text{Rep}(H)$ defined by a homomorphism $f: H \rightarrow G$ of affine group schemes. The objects of $\text{Rep}(G)^{\omega^f}$ are those on which H (equivalently, the intersection of the normal subgroups of G containing $f(H)$) acts trivially. The functor ω^f is a quotient functor if and only if f is a closed immersion, in which case it is normal if and only if $f(H)$ is normal in G .

PROPOSITION 2.2 *An exact tensor functor $q: \mathcal{T} \rightarrow \mathcal{Q}$ of tannakian categories is a normal quotient functor if and only if there exists a subgroup H of $\pi(\mathcal{T})$ such that $\pi(q)$ induces an isomorphism $\pi(\mathcal{Q}) \rightarrow q(H)$.*

PROOF. $\Leftarrow$: Because q is exact, $q(H) \rightarrow q(\pi\mathcal{T})$ is a closed immersion. Therefore $\pi(q)$ is a closed immersion, and its image is the normal subgroup $q(H)$ of $q(\pi\mathcal{T})$.

$\Rightarrow$: Because q is a quotient functor, $\pi(q)$ is a closed immersion. Let H be the kernel of the homomorphism $\pi(\mathcal{T}) \rightarrow \pi(\mathcal{T}^q)$ defined by the inclusion $\mathcal{T}^q \hookrightarrow \mathcal{T}$. The image of $\pi(q)$ is contained in $q(H)$, and equals it if and only if q is normal. To see this, let $G = q\pi(\mathcal{T})$, and identify $\mathcal{T}$ with the category of objects of $\mathcal{Q}$ with an action of G compatible with that

of $\pi(Q) \subset G$. Then q becomes the forgetful functor, and $T^q = T^{\pi(Q)}$. Thus, $q(H)$ is the subgroup of G acting trivially on those objects on which $\pi(Q)$ acts trivially. It follows that $\pi(Q) \subset q(H)$, with equality if and only if $\pi(Q)$ is normal in G . $\square$

In the situation of the proposition, we sometimes call Q a **quotient of T by H** (cf. Milne 2002, 1.3).

Let $q: T \rightarrow Q$ be an exact tensor functor of tannakian categories. By definition, q maps T^q into $Q^{\pi(Q)}$, and so we acquire a k -valued fibre functor $\omega^q \stackrel{\text{def}}{=} \gamma^Q \circ (q|_{T^q})$ on T^q :

$$\begin{array}{ccccc} & & \omega^q & & \\ & \nearrow & & \searrow & \\ T^q & \xrightarrow{q|_{T^q}} & Q^{\pi(Q)} & \xrightarrow{\gamma^Q} & \text{Vec}_k \\ \downarrow & & \downarrow & & \\ T & \xrightarrow{q} & Q. & & \end{array}$$

In particular, T^q is neutral. A fibre functor ω on Q , defines a fibre functor $\omega \circ q$ on T , and the (unique) isomorphism $\gamma^Q \rightarrow \omega|_{Q^{\pi(Q)}}$ defines an isomorphism $a(\omega): \omega^q \rightarrow (\omega \circ q)|_{T^q}$.

PROPOSITION 2.3 *Let $q: T \rightarrow Q$ be a normal quotient, and let H be the subgroup of $\pi(T)$ such that $\pi(Q) \simeq q(H)$.*

(a) *For X, Y in T , there is a canonical functorial isomorphism*

$$\text{Hom}_Q(qX, qY) \simeq \omega^q(\underline{\text{Hom}}(X, Y)^H).$$

(b) *The map $\omega \mapsto (\omega \circ q, a(\omega))$ defines an equivalence of gerbes*

$$r(q): \text{FIB}(Q) \rightarrow (\omega^q \downarrow \text{FIB}(T)).$$

PROOF. (a) From the various definitions and Deligne and Milne 1982,

$$\begin{aligned} \text{Hom}_Q(qX, qY) &\simeq \text{Hom}_Q(\mathbf{1}, \underline{\text{Hom}}(qX, qY)^{\pi(Q)}) && (\text{ibid. 1.6.4}) \\ &\simeq \text{Hom}_Q(\mathbf{1}, (q\underline{\text{Hom}}(X, Y))^{q(H)}) && (\text{ibid. 1.9}) \\ &\simeq \text{Hom}_Q(\mathbf{1}, q(\underline{\text{Hom}}(X, Y)^H)) \\ &\simeq \omega^q(\underline{\text{Hom}}(X, Y)^H) && (\text{definition of } \omega^q). \end{aligned}$$

(b) The functor $\text{FIB}(T) \rightarrow \text{FIB}(T^H)$ gives rise to an exact sequence

$$1 \rightarrow \text{Bd}(\omega_Q \downarrow \text{FIB}(T)) \rightarrow \text{Bd}(T) \rightarrow \text{Bd}(T^H) \rightarrow 0$$

(see 1.2). On the other hand, we saw in the proof of (2.2) that $H = \text{Ker}(\pi(T) \rightarrow \pi(T^H))$. On comparing these statements, we see that the morphism $r(q)$ of gerbes is bound by an isomorphism of bands, which implies that it is an equivalence of gerbs (Giraud 1971, IV 2.2.6). $\square$

PROPOSITION 2.4 *Let (Q, q) be a normal quotient of T . An exact tensor functor $q': T \rightarrow T'$ factors through q if and only if $T^{q'} \supset T^q$ and $\omega^q \approx \omega^{q'}|_{T^q}$.*

PROOF. The conditions are obviously necessary. For the sufficiency, choose an isomorphism $b: \omega^q \rightarrow \omega^{q'}|T^q$. A fibre functor ω on T' then defines a fibre functor $\omega \circ q'$ on T and an isomorphism $a(\omega)|T^q \circ b: \omega^q \rightarrow (\omega \circ q')|T^q$. In this way we get a homomorphism

$$\text{FIB}(T') \rightarrow (\omega^q \downarrow \text{FIB}(T)) \simeq \text{FIB}(Q)$$

and we can apply (1.4) to get a functor $Q \rightarrow T'$ with the correct properties. $\square$

THEOREM 2.5 *Let T be a tannakian category over k , and let ω_0 be a k -valued fibre functor on T^H for some subgroup $H \subset \pi(T)$. There exists a quotient (Q, q) of T by H such that $\omega^q \simeq \omega_0$.*

PROOF. The gerbe $(\omega_0 \downarrow \text{FIB}(T))$ is tannakian (see 1.3). From the morphism of gerbes

$$(\omega, a) \mapsto \omega: (\omega_0 \downarrow \text{FIB}(T)) \rightarrow \text{FIB}(T),$$

we obtain a morphism of tannakian categories

$$\text{Rep}(\text{FIB}(T)) \rightarrow \text{Rep}(\omega_0 \downarrow \text{FIB}(T))$$

(see 1.4). We define Q to be $\text{Rep}(\omega_0 \downarrow \text{FIB}(T))$ and we define q to be the composite of the above morphism with the equivalence (see 1.4)

$$T \rightarrow \text{Rep}(\text{FIB}(T)).$$

Since a gerbe and its tannakian category of representations have the same band, an argument as in the proof of Proposition 2.3 shows that $\pi(q)$ maps $\pi(Q)$ isomorphically onto $q(H)$. A direct calculation shows that ω^q is canonically isomorphic to ω_0 . $\square$

We sometimes write T/ω for the quotient of T defined by a k -valued fibre functor ω on a subcategory of T .

EXAMPLE 2.6 Let $(T, w, \mathbb{T})$ be a Tate triple, and let S be the full subcategory of T of objects isomorphic to a direct sum of integer tensor powers of the Tate object $\mathbb{T}$. Define ω_0 to be the fibre functor on S ,

$$X \mapsto \varinjlim_n \text{Hom}\left(\bigoplus_{-n \leq r \leq n} \mathbb{1}(r), X\right).$$

Then the quotient tannakian category T/ω_0 is that defined in Deligne and Milne 1982, 5.8.

REMARK 2.7 Let $q: T \rightarrow Q$ be a normal quotient functor. Then T can be recovered from Q , the homomorphism $\pi(Q) \rightarrow q(\pi(T))$, and the actions of $q(\pi(T))$ on the objects of Q (apply 1.6(a)).

REMARK 2.8 A fixed k -valued fibre functor on a tannakian category T determines a Galois correspondence between the subsets of $\text{ob}(T)$ and the equivalence classes of quotient functors $T \rightarrow Q$.

EXERCISE 2.9 Use (1.10, 1.11) to express the correspondence between fibre functors on tannakian subcategories of $\mathcal{T}$ and normal quotients of $\mathcal{T}$ in the language of 2-categories.

ASIDE 2.10 Let G be the fundamental group $\pi(\mathcal{T})$ of a tannakian category $\mathcal{T}$, and let H be a subgroup of G . We use the same letter to denote an affine group scheme in $\mathcal{T}$ and the band it defines. Then, under certain hypotheses, for example, if all the groups are commutative, there will be an exact sequence

$$\cdots \rightarrow H^1(k, G) \rightarrow H^1(k, G/H) \rightarrow H^2(k, H) \rightarrow H^2(k, G) \rightarrow H^2(k, G/H).$$

The category $\mathcal{T}$ defines a class $c(\mathcal{T})$ in $H^2(k, G)$, namely, the G -equivalence class of the gerbe of fibre functors on $\mathcal{T}$, and the image of $c(\mathcal{T})$ in $H^2(k, G/H)$ is the class of $\mathcal{T}^H$. Any quotient of $\mathcal{T}$ by H defines a class in $H^2(k, H)$ mapping to $c(\mathcal{T})$ in $H^2(k, G)$. Thus, the exact sequence suggests that a quotient of $\mathcal{T}$ by H will exist if and only if the cohomology class of $\mathcal{T}^H$ is neutral, i.e., if and only if $\mathcal{T}^H$ is neutral as a tannakian category, in which case the quotients are classified by the elements of $H^1(k, G/H)$ (modulo $H^1(k, G)$). When $\mathcal{T}$ is neutral and we fix a k -valued fibre functor on it, then the elements of $H^1(k, G/H)$ classify the k -valued fibre functors on $\mathcal{T}^H$. Thus, the cohomology theory suggests the above results, and in the next subsection we prove that a little more of this heuristic picture is correct.

The cohomology class of the quotient

For an affine group scheme G over a field k , $H^r(k, G)$ denotes the cohomology group computed with respect to the flat topology. When G is not commutative, this is defined only for $r = 0, 1, 2$ (Giraud 1971).

PROPOSITION 2.11 *Let (Q, q) be a quotient of $\mathcal{T}$ by a subgroup H of the centre of $\pi(\mathcal{T})$. Suppose that $\mathcal{T}$ is neutral, with k -valued fibre functor ω . Let $G = \underline{\text{Aut}}^\otimes(\omega)$, and let $\wp(\omega^q)$ be the $G/\omega(H)$ -torsor $\underline{\text{Hom}}(\omega|_{\mathcal{T}^H}, \omega^q)$. Under the connecting homomorphism*

$$H^1(k, G/H) \rightarrow H^2(k, H)$$

the class of $\wp(\omega^q)$ in $H^1(k, G/H)$ maps to the class of Q in $H^2(k, H)$.

PROOF. Note that $H = \text{Bd}(Q)$, and so the statement makes sense. According to Giraud 1971, IV 4.2.2, the connecting homomorphism sends the class of $\wp(\omega^q)$ to the class of the gerbe of liftings of $\wp(\omega^q)$, which can be identified with $(\omega^q \downarrow \text{FIB}(\mathcal{T}))$. Now Proposition 2.3 shows that the H -equivalence class of $(\omega^q \downarrow \text{FIB}(\mathcal{T}))$ equals that of $\text{FIB}(Q)$ which (by definition) is the cohomology class of Q . $\square$

Semisimple normal quotients

Everything can be made more explicit when the categories are semisimple. Throughout this subsection, k has characteristic zero.

PROPOSITION 2.12 *Every normal quotient of a semisimple tannakian category is semisimple.*

PROOF. A tannakian category is semisimple if and only if the identity component of its fundamental group is pro-reductive (cf. Deligne and Milne 1982, 2.28), and a normal subgroup of a reductive group is reductive (because its unipotent radical is a characteristic subgroup). $\square$

Let $\mathcal{T}$ be a semisimple tannakian category over k , and let ω_0 be a k -valued fibre functor on a tannakian subcategory S of $\mathcal{T}$. We can construct an explicit quotient $\mathcal{T}/\omega_0$ as follows. First, let $(\mathcal{T}/\omega_0)'$ be the category with one object $\overline{X}$ for each object X of $\mathcal{T}$, and with

$$\mathrm{Hom}_{(\mathcal{T}/\omega_0)'}(\overline{X}, \overline{Y}) = \omega_0(\underline{\mathrm{Hom}}(\overline{X}, \overline{Y})^H)$$

where H is the subgroup of $\pi(\mathcal{T})$ defining S . There is a unique structure of a k -linear tensor category on $(\mathcal{T}/\omega_0)'$ for which $q: \mathcal{T} \rightarrow (\mathcal{T}/\omega_0)'$ is a tensor functor. With this structure, $(\mathcal{T}/\omega_0)'$ is rigid, and we define $\mathcal{T}/\omega_0$ to be its pseudo-abelian hull. Thus, $\mathcal{T}/\omega_0$ has

$$\begin{aligned} \text{objects:} & \text{ pairs } (\overline{X}, e) \text{ with } X \in \mathrm{ob}(\mathcal{T}) \text{ and } e \text{ an idempotent in } \mathrm{End}(\overline{X}), \\ \text{morphisms:} & \mathrm{Hom}_{\mathcal{T}/\omega_0}((\overline{X}, e), (\overline{Y}, f)) = f \circ \mathrm{Hom}_{(\mathcal{T}/\omega_0)'}(\overline{X}, \overline{Y}) \circ e. \end{aligned}$$

Then $(\mathcal{T}/\omega_0, q)$ is a quotient of $\mathcal{T}$ by H , and $\omega^q \simeq \omega_0$.

Let ω be a fibre functor on $\mathcal{T}$, and let a be an isomorphism $\omega_0 \rightarrow \omega|_{\mathcal{T}^H}$. The pair (ω, a) defines a fibre functor ω_a on $\mathcal{T}/\omega_0$ whose action on objects is determined by

$$\omega_a(\overline{X}) = \omega(X)$$

and whose action on morphisms is determined by

$$\begin{array}{ccc} \mathrm{Hom}(\overline{X}, \overline{Y}) & \xrightarrow{\omega_a} & \mathrm{Hom}(\omega_a(\overline{X}), \omega_a(\overline{Y})) \\ \parallel \text{def} & & \uparrow \\ \omega_0(\underline{\mathrm{Hom}}(X, Y)^H) & \xrightarrow{a} \omega(\underline{\mathrm{Hom}}(X, Y)^H) \xrightarrow{\simeq} \underline{\mathrm{Hom}}(\omega(X), \omega(Y))^{\omega(H)} & \end{array}$$

The map $(\omega, a) \mapsto \omega_a$ defines an equivalence $(\omega_0 \downarrow \mathrm{FIB}(\mathcal{T})) \rightarrow \mathrm{FIB}(\mathcal{T}/\omega_0)$.

Let $H_1 \subset H_0 \subset \pi(\mathcal{T})$, and let ω_0 and ω_1 be k -valued fibre functors on $\mathcal{T}^{H_0}$ and $\mathcal{T}^{H_1}$ respectively. A morphism $\alpha: \omega_0 \rightarrow \omega_1|_{\mathcal{T}^{H_0}}$ defines an exact tensor functor $\mathcal{T}/\omega_0 \rightarrow \mathcal{T}/\omega_1$ whose action on objects is determined by

$$\overline{X} \text{ (in } \mathcal{T}^{H_0}) \mapsto \overline{X} \text{ (in } \mathcal{T}^{H_1}),$$

and whose action on morphisms is determined by

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{T}/\omega_0}(\overline{X}, \overline{Y}) & \xrightarrow{\quad} & \mathrm{Hom}_{\mathcal{T}/\omega_1}(\overline{X}, \overline{Y}) \\ \parallel \text{def} & & \parallel \text{def} \\ \omega_0(\underline{\mathrm{Hom}}_{\mathcal{T}}(X, Y)^{H_0}) & \xrightarrow{\alpha} \omega_1(\underline{\mathrm{Hom}}_{\mathcal{T}}(X, Y)^{H_0}) \hookrightarrow \omega_1(\underline{\mathrm{Hom}}_{\mathcal{T}}(X, Y)^{H_1}) & \end{array}$$

When $H_1 = H_0$, this is an isomorphism (!) of tensor categories $\mathbb{T}/\omega_0 \rightarrow \mathbb{T}/\omega_1$.

Let (Q_1, q_1) and (Q_2, q_2) be quotients of $\mathbb{T}$ by H . For simplicity, assume that $\pi \stackrel{\text{def}}{=} \pi(\mathbb{T})$ is commutative. Then $\underline{\text{Hom}}(\omega^{q_1}, \omega^{q_2})$ is π/H -torsor, and we assume that it lifts to a π -torsor P in $\mathbb{T}$, so $P \wedge^\pi (\pi/H) = \underline{\text{Hom}}(\omega^{q_1}, \omega^{q_2})$. Then

$$\mathbb{T} \xrightarrow{X \mapsto P \wedge^\pi X} \mathbb{T} \xrightarrow{q_2} Q_2$$

realizes Q_2 as a quotient of $\mathbb{T}$ by H , and the corresponding fibre functor on $\mathbb{T}^H$ is $P \wedge^\pi \omega^{q_2} \simeq \omega^{q_1}$. Therefore, there exists a commutative diagram of exact tensor functors

$$\begin{array}{ccc} \mathbb{T} & \xrightarrow{X \mapsto P \wedge^\pi X} & \mathbb{T} \\ \downarrow q_1 & & \downarrow q_2 \\ Q_1 & \longrightarrow & Q_2, \end{array}$$

which depends on the choice of P lifting $\underline{\text{Hom}}(\omega^{q_1}, \omega^{q_2})$ in an obvious way.

3 Polarizations

We refer to Deligne and Milne 1982, 5.12, for the notion of a (graded) polarization on a Tate triple over $\mathbb{R}$. We write $\mathbb{V}$ for the category of $\mathbb{Z}$ -graded complex vector spaces endowed with a semilinear automorphism a such that $a^2 v = (-1)^n v$ if $v \in V^n$. It has a natural structure of a Tate triple (ibid. 5.3). The canonical polarization on $\mathbb{V}$ is denoted $\Pi^\vee$.

A morphism $F: (\mathbb{T}_1, w_1, \mathbb{T}_1) \rightarrow (\mathbb{T}_2, w_2, \mathbb{T}_2)$ of Tate triples is an exact tensor functor $F: \mathbb{T}_1 \rightarrow \mathbb{T}_2$ preserving the gradations together with an isomorphism $F(\mathbb{T}_1) \simeq \mathbb{T}_2$. We say that such a morphism is **compatible** with graded polarizations Π_1 and Π_2 on $\mathbb{T}_1$ and $\mathbb{T}_2$ (denoted $F: \Pi_1 \mapsto \Pi_2$) if

$$\psi \in \Pi_1(X) \Rightarrow F\psi \in \Pi_2(FX),$$

in which case, for any X homogeneous of weight n , $\Pi_1(X)$ consists of the sesquilinear forms $\psi: X \otimes \overline{X} \rightarrow \mathbb{1}(-n)$ such that $F\psi \in \Pi_2(FX)$. In particular, given F and Π_2 , there exists at most one graded polarization Π_1 on $\mathbb{T}_1$ such that $F: \Pi_1 \mapsto \Pi_2$.

Let $\mathbb{T} = (\mathbb{T}, w, \mathbb{T})$ be an algebraic Tate triple over $\mathbb{R}$ such that $w(-1) \neq 1$. Given a graded polarization Π on $\mathbb{T}$, there exists a morphism of Tate triples $\xi_\Pi: \mathbb{T} \rightarrow \mathbb{V}$ (well defined up to isomorphism) such that $\xi_\Pi: \Pi \mapsto \Pi^\vee$ (Deligne and Milne 1982, 5.20). Let ω_Π be the composite

$$\mathbb{T}^{w(\mathbb{G}_m)} \xrightarrow{\xi_\Pi} \mathbb{V}^{w(\mathbb{G}_m)} \xrightarrow{\gamma^\vee} \text{Vec}_{\mathbb{R}};$$

it is a fibre functor on $\mathbb{T}^{w(\mathbb{G}_m)}$.

A criterion for the existence of a polarization

PROPOSITION 3.1 *Let $\mathbb{T} = (\mathbb{T}, w, \mathbb{T})$ be an algebraic Tate triple over $\mathbb{R}$ such that $w(-1) \neq 1$, and let $\xi: \mathbb{T} \rightarrow \mathbb{V}$ be a morphism of Tate triples. There exists a graded polarization Π on $\mathbb{T}$ (necessarily unique) such that $\xi: \Pi \mapsto \Pi^\vee$ if and only if the real algebraic group $\underline{\text{Aut}}^\otimes(\gamma^\vee \circ \xi|_{\mathbb{T}^{w(\mathbb{G}_m)}})$ is anisotropic.*

PROOF. Let $G = \underline{\text{Aut}}^{\otimes}(\gamma^V \circ \xi | T^{w(\mathbb{G}_m)})$. Assume Π exists. The restriction of Π to $T^{w(\mathbb{G}_m)}$ is a symmetric polarization, which the fibre functor $\gamma^V \circ \xi$ maps to the canonical polarization on $\text{Vec}_{\mathbb{R}}$. This implies that G is anisotropic (Deligne 1972, 2.6).

For the converse, let X be an object of weight n in $T_{(\mathbb{C})}$. A sesquilinear form $\psi: \xi(X) \otimes \overline{\xi(X)} \rightarrow \mathbf{1}(-n)$ arises from a sesquilinear form on X if and only if it is fixed by G . Because G is anisotropic, there exists a $\psi \in \Pi^V(\xi(X))$ fixed by G (ibid., 2.6), and we define $\Pi(X)$ to consist of all sesquilinear forms ϕ on X such that $\xi(\phi) \in \Pi^V(\xi(X))$. It is now straightforward to check that $X \mapsto \Pi(X)$ is a polarization on T . $\square$

COROLLARY 3.2 *Let $F: (T_1, w_1, \mathbb{T}_1) \rightarrow (T_2, w_2, \mathbb{T}_2)$ be a morphism of Tate triples, and let Π_2 be a graded polarization on T_2 . There exists a graded polarization Π_1 on T_1 such that $F: \Pi_1 \mapsto \Pi_2$ if and only if the real algebraic group $\underline{\text{Aut}}^{\otimes}(\gamma^V \circ \xi_{\Pi_2} \circ F | T_1^{w(\mathbb{G}_m)})$ is anisotropic.*

Polarizations on quotients

The next proposition gives a criterion for a polarization on a Tate triple to pass to a quotient Tate triple.

PROPOSITION 3.3 *Let $T = (T, w, \mathbb{T})$ be an algebraic Tate triple over $\mathbb{R}$ such that $w(-1) \neq 1$. Let (Q, q) be a quotient of T by $H \subset \pi(T)$, and let ω^q be the corresponding fibre functor on T^H . Assume $H \supset w(\mathbb{G}_m)$, so that Q inherits a Tate triple structure from that on T , and that Q is semisimple. Given a graded polarization Π on T , there exists a graded polarization Π' on Q such that $q: \Pi \mapsto \Pi'$ if and only if $\omega^q \approx \omega_{\Pi} | T^H$.*

PROOF. $\Rightarrow$: Let Π' be such a polarization on Q , and consider the functors

$$T \xrightarrow{q} Q \xrightarrow{\xi_{\Pi'}} V, \quad \xi_{\Pi'}: \Pi' \mapsto \Pi^V.$$

Both $\xi_{\Pi'} \circ q$ and ξ_{Π} are compatible with Π and Π^V and with the Tate triple structures on T and V , and so $\xi_{\Pi'} \circ q \approx \xi_{\Pi}$ (Deligne and Milne 1982, 5.20). On restricting everything to $T^{w(\mathbb{G}_m)}$ and composing with γ^V , we get an isomorphism $\omega_{\Pi'} \circ (q | T^{w(\mathbb{G}_m)}) \approx \omega_{\Pi}$. Now restrict this to T^H , and note that

$$(\omega_{\Pi'} \circ (q | T^{w(\mathbb{G}_m)})) | T^H = (\omega_{\Pi'} | Q^{\pi(Q)}) \circ (q | T^H) \simeq \omega^q$$

because $\omega_{\Pi'} | Q^{\pi(Q)} \simeq \gamma^Q$.

$\Leftarrow$: The choice of an isomorphism $\omega^q \rightarrow \omega_{\Pi} | T^H$ determines an exact tensor functor

$$T/\omega^q \rightarrow T/\omega_{\Pi}.$$

As the quotients T/ω^q and T/ω_{Π} are tensor equivalent respectively to Q and V , this shows that there is an exact tensor functor $\xi: Q \rightarrow V$ such that $\xi \circ q \approx \xi_{\Pi}$. Evidently $\underline{\text{Aut}}^{\otimes}(\gamma^V \circ \xi | Q^{w(\mathbb{G}_m)})$ is isomorphic to a subgroup of $\underline{\text{Aut}}^{\otimes}(\gamma^V \circ \xi_{\Pi} | T^{w(\mathbb{G}_m)})$. Since the latter is anisotropic, so also is the former (Deligne 1972, 2.5). Hence ξ defines a graded polarization Π' on Q (Proposition 3.1), and clearly $q: \Pi \mapsto \Pi'$. $\square$

References

- BERTOLIN, C. 2003. Motivic Galois theory for motives of niveau ≤ 1 . Preprint arXiv:math.NT/0309379.
- DELIGNE, P. 1972. La conjecture de Weil pour les surfaces $K3$. *Invent. Math.* 15:206–226.
- DELIGNE, P. 1989. Le groupe fondamental de la droite projective moins trois points, pp. 79–297. *In* Galois groups over $\mathbf{Q}$ (Berkeley, CA, 1987), volume 16 of *Math. Sci. Res. Inst. Publ.* Springer, New York.
- DELIGNE, P. 1990. Catégories tannakiennes, pp. 111–195. *In* The Grothendieck Festschrift, Vol. II, Progr. Math. Birkhäuser Boston, Boston, MA.
- DELIGNE, P. AND MILNE, J. S. 1982. Tannakian categories, pp. 101–228. *In* Hodge cycles, motives, and Shimura varieties, Lecture Notes in Mathematics. Springer-Verlag, Berlin.
- GIRAUD, J. 1971. Cohomologie non abélienne. Springer-Verlag, Berlin.
- MILNE, J. S. 2002. Polarizations and Grothendieck’s standard conjectures. *Ann. of Math. (2)* 155:599–610.
- MILNE, J. S. 2007. Rational Tate classes on abelian varieties. In preparation.
- SAAVEDRA RIVANO, N. 1972. Catégories Tannakiennes. Springer-Verlag, Berlin.
- J.S. Milne, Ann Arbor, MI, USA